

JOURNAL OF FUNCTIONAL ANALYSIS **101**, 25–37 (1991)

On Cauchy–Kovalewski Extension

G. LAVILLE

*Université de Caen, Département de Mathématiques,
14032 Caen Cedex, France**Communicated by P. Malliavin*

Received April 1990

Let \mathcal{A} be the universal Clifford algebra of the euclidean space \mathbb{R}^m , e_1, \dots, e_m the euclidean basis, $D = \partial/\partial x_0 + \sum_{i=1}^m e_i \partial/\partial x_i$ the main operator of Clifford analysis. Given $f: \mathbb{R}^m \rightarrow \mathbb{R}$ a certain kind of rational function, it is possible to find explicitly the extension of f , $f^*: \mathbb{R}^{m+1} \rightarrow \mathcal{A}$, satisfying $Df^* = 0$. This also gives the harmonic extension of f . f^* is given by integral representation on simplex in \mathbb{C}^n . © 1991 Academic Press, Inc.

INTRODUCTION

In [1], the Cauchy–Kovalewski extension was defined. It is known how to extend polynomials and hence how to extend real analytic functions.

Here, we show how to extend rational functions which have a factorization in the product of functions of first degree.

The motivations of that problem are as follows:

- there are no good methods of harmonic extension of rational functions
- rational functions are very important in the theory of one complex variable. What are their roles in Clifford analysis?
- the Dirac equation with an external electromagnetic field would be better understood if we knew how to manage multiple singularities.

NOTATION

Recall some notations from [1]. \mathcal{A} is the universal Clifford algebra of the euclidean space \mathbb{R}^m but with the minus sign for the scalar product. e_1, \dots, e_m is the euclidean basis and

$$D = \frac{\partial}{\partial x_0} + \sum_{i=1}^m e_i \frac{\partial}{\partial x_i}$$

is the generalized Cauchy–Riemann operator acting on functions defined on the \mathbb{R}^{m+1} space. A function is (left) monogenic when $Df=0$,

Let $\Omega \subset \mathbb{R}^m$ be open and $f: \Omega \rightarrow \mathbb{R}$.

The maximal (left) monogenic extension $f^*: U \rightarrow \mathcal{A}$, U open in \mathbb{R}^{m+1} , $f^*=f$ on Ω , $Df^*=0$, is called the (left) Cauchy–Kovalewski extension.

\mathcal{A} is thought to be imbedded in his complexified $\mathcal{A} \otimes_{\mathbb{R}} \mathbb{C}$.

The standard p -simplex is denoted by

$$\Sigma_p = \left\{ t^{(p)} \in \mathbb{R}^p; \sum_{j=1}^p t_j \leq 1 \text{ and } 0 \leq t_j \text{ for all } j \right\};$$

elements of \mathbb{R}^p are $t^{(p)} = (t_1, \dots, t_p)$,

$$A_{p-1} = \left\{ t^{(p)} \in \mathbb{R}^p; \sum_{j=1}^p t_j = 1 \text{ and } 0 \leq t_j \text{ for all } j \right\}.$$

Let $v_1, \dots, v_k \in \mathcal{A}$ and $z = (z_1, \dots, z_k) \in \mathbb{C}^k$; then $\langle z, v \rangle = \sum_{j=1}^k z_j v_j$,

$$\omega(z) = dz_1 \wedge \dots \wedge dz_k$$

$$\omega'(z) = \sum_{l=1}^k (-1)^{l-1} z_l dz_1 \wedge \dots \wedge \widehat{dz_l} \wedge \dots \wedge dz_k.$$

Note that $d\omega'(z) = k\omega(z)$.

Let $j = (j_1, \dots, j_k) \in \mathbb{N}^k$ a multiindice; then $|j| = j_1 + \dots + j_k$, $z^j = z_1^{j_1} \dots z_k^{j_k}$, $j! = j_1! \dots j_k!$. Let $\alpha = (\alpha_1, \dots, \alpha_k) \in \mathbb{N}^k$; then $D_{\zeta,0}^\alpha = \partial^{x_1+\dots+x_k}/\partial \zeta_1^{\alpha_1} \dots \partial \zeta_k^{\alpha_k}$ is taken for $\zeta=0$.

If E is a set, $\forall E$ is a neighborhood of E .

1. HOMOTOPY AND CLIFFORD-VALUED INTEGRALS

In this section, we define, study, and compute some Clifford valued integrals based on deformations of simplex.

LEMMA 1. *Let $a, b, c \in \mathcal{A}$ and $p, q \in \mathbb{N}$ such that $p \geq q$ and $v \subset \mathbb{C}$ a closed path such that $a + zb + (1-z)c$ is invertible for all $z \in v$. Then*

$$\int_v \frac{(1-z)^q dz}{[a + zb + (1-z)c]^{p+2}} = 0.$$

Proof. The Clifford algebra \mathcal{A} may be thought of as a matrix algebra; then the singularities of $1/[a + zb + (1-z)c]^{p+2}$ are isolated. Then, we

must show only that for all singularities z_0 there exists a neighborhood $V\{z_0\}$ of z_0 such that

$$\int_v \frac{(1-z)^q dz}{[a+zb+(1-z)c]^{p+2}} = 0$$

with $v \subset V\{z_0\}$.

The first case is for $b-c$ invertible:

$$\begin{aligned} & \frac{1}{a+zb+(1-z)c} \\ &= \frac{1}{(z-z_0)(b-c)+z_0(b-c)+a+c} \\ &= \sum_{k=0}^{\infty} (-1)^k \frac{1}{(z-z_0)^{k+1}} (b-c)^{-1} [z_0 + (b-c)^{-1}(a+c)]^k. \end{aligned}$$

We can take $V\{z_0\}$ such that this series is uniformly convergent on $V\{z_0\}$,

$$\begin{aligned} & \int_v \frac{(1-z)^q dz}{[a+zb+(1-z)c]^{p+2}} \\ &= \int_v \left[\sum_{k=0}^{\infty} (-1)^k \frac{1-z_0-(z-z_0)}{(z-z_0)^{k+1}} (b-c)^{-1} (z_0 + (b-c)^{-1}(a+c))^k \right]^q \\ & \quad \times \left[\sum_{h=0}^{\infty} (-1)^h \frac{1}{(z-z_0)^{h+1}} (b-c)^{-1} (z_0 + (b-c)^{-1}(a+c))^h \right]^{p-q+2} dz = 0 \end{aligned}$$

because we have no term involving $1/(z-z_0)$. ■

PROPOSITION 1. *Let m_1, \dots, m_k , k invertible elements of the Clifford algebra \mathcal{A} . There exists C^∞ mapping*

$$\gamma_p: V\Sigma_p \rightarrow \mathbb{C}, \quad 1 \leq p \leq k-1,$$

such that

$$\gamma_p(1, 0, \dots, 0) = 1;$$

$$\gamma_p(0, t_2, \dots, t_p) = 0 \quad \text{for all } (t_2, \dots, t_p) \in \Sigma_{p-1}$$

$$\Gamma: V\Sigma_{k-1} \rightarrow \mathbb{C}^{k-1}$$

$$t^{(k-1)} \mapsto (\gamma_{k-1}(t^{(k-1)}), \dots, \gamma_1(t^{(1)}))$$

$z_1 m_1 + (1 - z_1)(z_2 m_2 + (1 - z_2)(\dots (z_{k-1} m_{k-1} + (1 - z_{k-1}) m_k) \dots)$ is invertible for all $z_p \in \gamma_p(t^{(p)})$. The integral

$$I = \int_{\Gamma(\Sigma_{k-1})} \frac{(1 - z_1)^{k-2} (1 - z_2)^{k-3} \dots (1 - z_{k-2}) dz_1 \wedge \dots \wedge dz_{k-1}}{\{z_1 m_1 + (1 - z_1)(z_2 m_2 + (1 - z_2)(\dots (z_{k-1} m_{k-1} + (1 - z_{k-1}) m_k) \dots)\}^k}$$

does not depend on the particular choice of the maps $\gamma_1, \dots, \gamma_{k-1}$.

Proof. Once again we use the isolated singularity property: there exists a path

$$\gamma_1: V \Sigma_1 \rightarrow \mathbb{C}, \quad \gamma_1(0) = 0, \quad \gamma_1(1) = 1,$$

such that $\zeta m_1 + (1 - \zeta) m_2$ is invertible on this path. Hence, the integral

$$\int_{\gamma_1(\Sigma_1)} \frac{d\zeta}{[\zeta m_1 + (1 - \zeta) m_2]^2} = \int_{\Sigma_1} \frac{\gamma_1'(t) dt}{[\gamma_1(t) m_1 + (1 - \gamma_1(t)) m_2]^2}$$

is well defined, and by Lemma 1 does not depend on the map γ_1 .

This proposition is true for $k = 1$. Suppose that it is also true for $k - 1$: Fix $t^{(k-1)} \in \Sigma_{k-1}$; there exists a path

$$\gamma_k(\cdot, t^{(k-1)}): V[(0, t^{(k-1)}), (1, 0)] \rightarrow \mathbb{C}$$

such that

$$\gamma_k(0, t^{(k-1)}) = 0, \quad \forall t^{(k-1)} \in \Sigma_{k-1}$$

$$\gamma_k(1, 0) = 1$$

and, letting

$$\begin{aligned} m(t^{(k-1)}) &= \gamma_{k-1}(t^{(k-1)}) m_1 + (1 - \gamma_{k-1}(t^{(k-1)})) (\gamma_{k-2}(t^{(k-2)}) m_2 \\ &\quad + \dots + (1 - \gamma_1(t^{(1)})) m_k) \dots), \end{aligned}$$

$\zeta m_0 + (1 - \zeta) m(t^{(k-1)})$ is invertible on that path.

The following integral is well defined:

$$\begin{aligned} I &= \int_{\gamma_1(\Sigma_1)} \int_{\gamma_2(\Sigma_2)} \dots \\ &\quad \dots \int_{\gamma_{k-1}(\Sigma_{k-1})} \frac{(1 - z_0)^{k-1} dz_0 \wedge (1 - z_1)^{k-2} \dots (1 - z_{k-2}) dz_1 \wedge \dots \wedge dz_{k-1}}{[z_0 m_0 + (1 - z_0)(\dots)]^{k+1}}. \end{aligned}$$

By induction this integral does not depend on the particular path $\gamma_2, \dots, \gamma_{k-1}$ and by Lemma 1 it does not depend on the particular path γ_1 . ■

PROPOSITION 2. *There exists a \mathcal{C}^∞ map $\psi: V\Sigma_{k-1} \rightarrow \mathbb{C}^{k-1}$ such that*

$$\psi(0, \dots, 0) = (0, \dots, 0); \quad \psi(0, \dots, 0, 1, 0, \dots, 0) = (0, \dots, 0, 1, 0, \dots, 0);$$

$\zeta \in \psi(\Sigma_{k-1}) \Rightarrow \sum_{j=1}^{k-1} \zeta_j m_j + (1 - \sum_{j=1}^{k-1} \zeta_j) m_k$ invertible. The integral of Proposition 1 is

$$I = \int_{\psi(\Sigma_{k-1})} \frac{d\zeta_1 \wedge \dots \wedge d\zeta_{k-1}}{\left[\sum_{j=1}^{k-1} \zeta_j m_j + \left(1 - \sum_{j=1}^{k-1} \zeta_j \right) m_k \right]^k}.$$

Proof. Change the variables in Proposition 1:

$$\begin{aligned} \zeta_1 &= z_1 \\ \zeta_2 &= (1 - z_1)z_2 \\ &\vdots \\ \zeta_{k-1} &= (1 - z_1)(1 - z_2) \cdots (1 - z_{k-2})z_{k-1}. \end{aligned}$$

Then

$$\begin{aligned} d\zeta_1 \wedge \dots \wedge d\zeta_{k-1} &= dz_1 \wedge ((1 - z_1) dz_2 - z_2 dz_1) \wedge \dots \\ &\quad \wedge \left(\sum_{j=1}^{k-2} (1 - z_1) \cdots \widehat{(1 - z_j)} \cdots (1 - z_{k-2}) z_{k-1} (-1) dz_j \right. \\ &\quad \left. + (1 - z_1) \cdots (1 - z_{k-2}) dz_{k-1} \right) \\ &= (1 - z_1)^{k-2} \cdots (1 - z_{k-2}) dz_1 \wedge \dots \wedge dz_{k-1}. \end{aligned}$$

The coefficient of m_1 is $z_1 = \zeta_1$, m_2 is $(1 - z_1)z_2 = \zeta_2$, and so on, ..., of m_{k-1} is $(1 - z_1)(1 - z_2) \cdots (1 - z_{k-2})z_{k-1} = \zeta_{k-1}$, of m_k is $(1 - z_1) \cdots (1 - z_{k-2})(1 - z_{k-1}) = 1 - \zeta_1 - \zeta_2 - \cdots - \zeta_{k-1}$.

Composing the map Γ of Proposition 1 and this change of variables, we get a map

$$\begin{aligned} V\Sigma_{k-1} &\rightarrow \mathbb{C}^{k-1} \\ t^{(k-1)} &\mapsto (\gamma_{k-1}(t^{(k-1)}), \dots, [1 - \gamma_{k-1}(t^{(k-1)})] \cdots [1 - \gamma_2(t^{(2)})] \gamma_1(t^{(1)})) \end{aligned}$$

which is a \mathcal{C}^∞ mapping such that

$$\begin{aligned} (0, \dots, 0) &\mapsto (0, \dots, 0) \\ (0, \dots, 0, 1, 0, \dots, 0) &\mapsto (0, \dots, 0, 1, 0, \dots, 0) \end{aligned}$$

and $\sum_{j=1}^{k-1} \zeta_j m_j + (1 - \sum_{j=1}^{k-1} \zeta_j) m_k$ is invertible for ζ belonging to the image of that map. Let ψ be this map. ■

THEOREM 1. *Let m_1, \dots, m_k , k invertible elements of the Clifford algebra \mathcal{A} . There exists a simplex $A \subset \mathbb{C}^k$, defined by the image of A_{k-1} by a \mathcal{C}^∞ mapping such that*

(i) $z \in A \Rightarrow \langle z, m \rangle$ is invertible in \mathcal{A}

(ii) $I = (-1)^{k-1} \int_A \omega'(z) / \langle z, m \rangle^k$ does not depend on the particular choice of A .

Proof. Consider two changes of variables:

$$f: \mathbb{C}^{k-1} \rightarrow \mathbb{C}^k$$

$$\zeta \mapsto z$$

$$z_1 = \zeta_1, \dots, z_{k-1} = \zeta_{k-1}, \quad z_k = 1 - \sum_{j=1}^{k-1} \zeta_j$$

$$\tilde{f}: \Sigma_{k-1} \rightarrow A_{k-1}$$

$$t \mapsto x$$

$$x_1 = t_1, \dots, x_{k-1} = t_{k-1}, \quad x_k = 1 - \sum_{j=1}^{k-1} t_j.$$

Let $\tilde{\psi} = \phi \circ \psi \circ \tilde{f}^{-1}$. We have

$$\sum_{p=1}^k (-1)^{p-1} z_p dz_1 \wedge \dots \wedge \widehat{dz_p} \wedge \dots \wedge dz_k = (-1)^{k-1} d\zeta_1 \wedge \dots \wedge d\zeta_{k-1}.$$

Hence, the integral I of Proposition 2 is equal to

$$I = (-1)^{k-1} \int_{\tilde{\psi}(A_{k-1})} \frac{\omega'(z)}{\langle z, m \rangle^k}.$$

We can put $A = \tilde{\psi}(A_{k-1})$.

The integral I does not depend on the particular choice of A because it is possible to proceed in the opposite way; starting with a A we make the inverse change of variables, getting the integral of proposition 2 and the integral of Proposition 1 which do not depend on the particular path. ■

PROPOSITION 3. *Let a_1, \dots, a_k , k strictly positive real numbers. Then*

$$(-1)^{k-1} (k-1)! \int_{A_{k-1}} \frac{\omega'(x)}{\langle x, a \rangle^k} = \frac{1}{a_1 \dots a_k}.$$

Proof. The differential form $\omega'(x)/\langle x, a \rangle^k$ is closed:

$$\begin{aligned} d \frac{\omega'(x)}{\langle x, a \rangle^k} &= k \frac{dx_1 \wedge \cdots \wedge dx_k}{\langle x, a \rangle^k} - k \frac{1}{\langle x, a \rangle^{k+1}} d\langle x, a \rangle \wedge \omega'(x) \\ &= k \frac{dx_1 \wedge \cdots \wedge dx_k}{\langle x, a \rangle^k} - k \frac{1}{\langle x, a \rangle^{k+1}} \sum_{j=1}^k a_j dx_j \wedge \omega'(x) \\ &= k \frac{dx_1 \wedge \cdots \wedge dx_k}{\langle x, a \rangle^k} - k \frac{1}{\langle x, a \rangle^{k+1}} \sum_{j=1}^k a_j x_j \omega(x) \\ &= 0. \end{aligned}$$

Let

$$\begin{aligned} B_{k-1} &= \left\{ x \in \mathbb{R}^k; \sum_{j=1}^k a_j x_j = 1, 0 \leq x_j \right\} \\ \Omega &= \{x \in \mathbb{R}^k; x = t\alpha + (1-t)\beta, t \in [0, 1], \alpha \in A_{k-1}, \beta \in B_{k-1}\} \\ C_j &= \{x \in \mathbb{R}^k; x_j = 0\} \cap \Omega. \end{aligned}$$

The boundary of Ω is $A_{k-1} \cup B_{k-1} \cup \bigcup_j C_j$. On C_j , $x_j = 0$; hence $\omega'(x) = 0$:

$$0 = \int_{\Omega} d \frac{\omega'(x)}{\langle x, a \rangle^k} = - \int_{A_{k-1}} \frac{\omega'(x)}{\langle x, a \rangle^k} + \int_{B_{k-1}} \frac{\omega'(x)}{\langle x, a \rangle^k} - \sum_{j=1}^k \int_{C_j} \frac{\omega'(x)}{\langle x, a \rangle^k}.$$

Hence

$$\int_{A_{k-1}} \frac{\omega'(x)}{\langle x, a \rangle^k} = \int_{B_{k-1}} \frac{\omega'(x)}{\langle x, a \rangle^k}.$$

We set

$$\begin{aligned} y_j &= a_j x_j \Leftrightarrow x_j = \frac{y_j}{a_j} \\ \int_{B_{k-1}} \frac{\omega'(x)}{\langle x, a \rangle^k} &= \int_{A_{k-1}} \frac{\omega'(y/a)}{\sum_{j=1}^k y_j} = \frac{1}{a_1 \cdots a_k} \int_{A_{k-1}} \omega'(y) \\ \int_{A_{k-1}} \omega'(y) &= (-1)^{k-1} \int_{\Sigma_{k-1}} dy_1 \wedge \cdots \wedge dy_{k-1} = \frac{(-1)^{k-1}}{(k-1)!}. \quad \blacksquare \end{aligned}$$

PROPOSITION 4. Let $a_1, \dots, a_k \in \mathbb{C} \setminus \{0\}$ and $\Lambda \subset \mathbb{C}^k$ a simplex image of A_{k-1} by a \mathcal{C}^∞ mapping such that $z \in \Lambda \Rightarrow \langle z, a \rangle \neq 0$. Then

$$(-1)^{k-1} (k-1)! \int_{\Lambda} \frac{\omega'(z)}{\langle z, a \rangle^k} = \frac{1}{a_1 \cdots a_k}.$$

Proof. Theorem 1 prove that such a simplex Λ exists. We must show the equality. Let

$$F = \{a \in \mathbb{C}^k; \operatorname{Re} a_j > 0, \operatorname{Im} a_j = 0 \text{ for all } 1 \leq j \leq k\}.$$

Let E be the connected component of the set $\{a \in (\mathbb{C} \setminus \{0\})^k$; there exist Λ , $z \in \Lambda \Rightarrow \langle z, a \rangle \neq 0$ and $(-1)^{k-1}(k-1)! \int_{\Lambda} \omega'(z) / \langle z, a \rangle^k = 1/a_1 \cdots a_k\}$ containing F : $F \subset E$ (by Proposition 3), $E \neq \emptyset$.

E is closed in $(\mathbb{C} \setminus \{0\})^k$ because for all sequences $(a_n) \rightarrow a$ with

$$(-1)^{k-1}(k-1)! \int_{\Lambda_n} \frac{\omega'(z)}{\langle z, a_n \rangle^k} = \frac{1}{a_{1,n} \cdots a_{k,n}},$$

if n is sufficiently great, we can take $\Lambda_n = \Lambda$ with Λ such that $z \in \Lambda \Rightarrow \langle z, a \rangle \neq 0$ and we can pass to the limit in the last equality.

E is open in $(\mathbb{C} \setminus \{0\})^k$ because if $\forall z \in \Lambda$, $\langle z, a \rangle \neq 0$, it is also true in a neighborhood of a . Moreover, the holomorphic functions of a ,

$$(-1)^{k-1}(k-1)! \int_{\Lambda} \frac{\omega'(z)}{\langle z, a \rangle^k} \text{ and } \frac{1}{a_1 \cdots a_k},$$

are equal in F and then are equal on the set E which is open.

Finally $E = (\mathbb{C} \setminus \{0\})^k$. ■

PROPOSITION 5. Let a_1, \dots, a_k and Λ be as in Proposition 4. Then

$$\frac{(-1)^{k-1}(|j|-1)!}{(j-1)!} \int_{\Lambda} \frac{z^{j-1} \omega'(z)}{\langle z, a \rangle^{|j|}} = \frac{1}{a_1^{j_1} \cdots a_k^{j_k}}.$$

Proof. Derivation with respect to the parameters a_1, \dots, a_k . ■

THEOREM 2. Let $a_1, \dots, a_k \in \mathbb{C} \setminus \{0\}$, $b_1, \dots, b_v \in \mathbb{C}$, Λ defined as in Proposition 4, α and j two multiindices such that $|j| > |\alpha|$. Then

$$C_{j,\alpha} D_{\zeta,0}^{\alpha} \int_{\Lambda} \frac{z^{j-1} \omega'(z)}{[\langle \zeta, b \rangle + \langle z, a \rangle]^{|j|-|\alpha|}} = \frac{b_1^{\alpha_1} \cdots b_v^{\alpha_v}}{a_1^{j_1} \cdots a_k^{j_k}}$$

with $C_{j,\alpha} = (-1)^{|\alpha|+k-1}(|j|-|\alpha|-1)!/(j-1)!$.

Proof. The integral is well defined, because if $\langle z, a \rangle \neq 0$ then for ζ sufficiently small $\langle \zeta, b \rangle + \langle z, a \rangle \neq 0$ (ζ small uniformly with respect to $z \in \Lambda$). The equality is proved by derivation under the integral sign. ■

Remark. The hypothesis $|j| > |\alpha|$ is not an actual restriction because it is possible to introduce an extra $a_{k+1} = 1$ and an integer j_{k+1} such that $|j| + j_{k+1} > |\alpha|$.

2. CAUCHY-KOVALEWSKI EXTENSION

For any function f defined on an open set of \mathbb{R}^m we denote by f^* its Cauchy-Kovalewski extension.

PROPOSITION 6. *Let $a_1, \dots, a_k, b_1, \dots, b_r$ be linear affine functions defined on \mathbb{R}^m with complex values, each non-identically zero. Let j and α be two multiindices such that $|j| > |\alpha|$ and*

$$U = \{(x_0, x) \in \mathbb{R}^{m+1}; a_l^*(x_0, x) \text{ invertible for all } 1 \leq l \leq k\}.$$

The Cauchy-Kovalewski extension of

$$\frac{b_1(x)^{x_1} \cdots b_r(x)^{x_r}}{a_1(x)^{j_1} \cdots a_k(x)^{j_k}}$$

is given for all $(x_0, x) \in U$ by

$$C_{j,\alpha} D_{\zeta,0}^\alpha \int_{A(x_0,x)} \frac{z^{j-1} \omega'(z)}{[\langle \zeta, b^*(x_0, x) \rangle + \langle z, a^*(x_0, x) \rangle]^{|j|-|\alpha|}}.$$

$A(x_0, x)$ is a simplex of type defined in Theorem 1. It is possible to choose the simplex $A(x_0, x)$ locally independent of the point (x_0, x) . The integral does not depend on the particular choice of that simplex.

Proof. $\partial a_l(x)/\partial x_i$ and $\partial b_l(x)/\partial x_i$ are constant; then their Cauchy-Kovalewski extensions are given by

$$\begin{aligned} a_l^*(x_0, x) &= a_l(x) - x_0 \sum_{i=1}^m e_i \frac{\partial a_l(x)}{\partial x_i} \\ b_l^*(x_0, x) &= b_l(x) - x_0 \sum_{i=1}^m e_i \frac{\partial b_l(x)}{\partial x_i}. \end{aligned}$$

By Theorem 1, there exists a simplex $A(x_0, x)$ such that the integral is well defined for all $(x_0, x) \in U$, and does not depend on the choice of that simplex: for all $(y_0, y) \in U$ there exists a neighborhood $V\{(y_0, y)\} \subset U$ such that if $(x_0, x) \in V\{(y_0, y)\}$ then

$$\begin{aligned} & C_{j,\alpha} D_{\zeta,0}^\alpha \int_{A(x_0,x)} \frac{z^{j-1} \omega'(z)}{[\langle \zeta, b^*(x_0, x) \rangle + \langle z, a^*(x_0, x) \rangle]^{|j|-|\alpha|}} \\ &= C_{j,\alpha} D_{\zeta,0}^\alpha \int_{A(y_0,y)} \frac{z^{j-1} \omega'(z)}{[\langle \zeta, b^*(x_0, x) \rangle + \langle z, a^*(x_0, x) \rangle]^{|j|-|\alpha|}}. \end{aligned}$$

We get a monogenic function because

$$\begin{aligned}
 & \langle \zeta, b^*(x_0, x) \rangle + \langle z, a^*(x_0, x) \rangle \\
 &= \sum_{l=1}^r \zeta_l \left(b_l(x) - x_0 \sum_{i=1}^m e_i \frac{\partial b_l(x)}{\partial x_i} \right) + \sum_{l=1}^k z_l \left(a_l(x) - x_0 \sum_{i=1}^m e_i \frac{\partial a_l(x)}{\partial x_i} \right) \\
 &= \left[\sum_{l=1}^r \zeta_l b_l(x) + \sum_{l=1}^k z_l a_l(x) \right] - x_0 \left[\sum_{i=1}^m e_i \left(\sum_{l=1}^r \zeta_l \frac{\partial b_l(x)}{\partial x_i} + \sum_{l=1}^k \frac{\partial a_l(x)}{\partial x_i} \right) \right] \\
 &= \varphi(x) - x_0 m \quad \text{with } m \in \mathcal{A}.
 \end{aligned}$$

Hence,

$$\begin{aligned}
 D \frac{1}{(\varphi(x) - x_0 m)^{|j| - |z|}} &= (|\alpha| - |j|) D(\varphi(x) - x_0 m) \frac{1}{(\varphi(x) - x_0 m)^{|j| - |\alpha| + 1}} \\
 &= 0.
 \end{aligned}$$

For the boundary values on \mathbb{R}^m , we have the following. Let $(0, y) \in U \cap \mathbb{R}^m$,

$$\begin{aligned}
 & \lim_{(x_0, x) \rightarrow (0, y)} C_{j, z} D_{\zeta, 0}^\alpha \int_{A(0, y)} \frac{z^{j-1} \omega'(z)}{[\langle \zeta, b^*(x_0, x) \rangle + \langle z, a^*(x_0, x) \rangle]^{|j| - |z|}} \\
 &= C_{j, a} D_{\zeta, 0}^\alpha \int_{A(0, y)} \frac{z^{j-1} \omega'(z)}{[\langle \zeta, b(x) \rangle + \langle z, a(x) \rangle]^{|j| - |z|}} \\
 &= \frac{b_1(x)^{x_1} \dots b_r(x)^{x_r}}{a_1(x)^{j_1} \dots a_k(x)^{j_k}}. \quad \blacksquare
 \end{aligned}$$

THEOREM 3. Let $a_1, \dots, a_k, b_1, \dots, b_r$ be linear affine functions: $\mathbb{R}^m \rightarrow \mathbb{R}$ each non-identically zero. Let j and α be two multiindices such that $|j| > |\alpha|$ and

$$\begin{aligned}
 U &= \{(x_0, x) \in \mathbb{R}^{m+1}; x_0 \neq 0\} \\
 &\cup \{(x_0, x) \in \mathbb{R}^{m+1}; x_0 = 0 \text{ and } a_l(x) \neq 0 \text{ for all } 1 \leq l \leq k\}.
 \end{aligned}$$

The Cauchy-Kovalewski extension of

$$\frac{b_1(x)^{x_1} \dots b_r(x)^{x_r}}{a_1(x)^{j_1} \dots a_k(x)^{j_k}}$$

is given for all $(x_0, x) \in U$ by

$$C_{j, a} D_{\zeta, 0}^\alpha \int_{A(x_0, x)} \frac{z^{j-1} \omega'(z)}{[\langle \zeta, b^*(x_0, x) \rangle + \langle z, a^*(x_0, x) \rangle]^{|j| - |z|}}.$$

The harmonic extension of the previous rational function defined on \mathbb{R}^m is the real part of that Clifford-valued function.

Proof. By Proposition 6 we must study only the set of all (x_0, x) such that $a^*(x_0, x)$ is invertible. Namely, we prove that, for all (x_0, x) such that $x_0 \neq 0$, $a_l^*(x_0, x)$ is invertible:

$$\frac{\partial a_l(x)}{\partial x_i} \text{ is constant.}$$

If, for all i , this constant is 0, then $a_l(x)$ is a constant not zero (because a_l is non-identically 0) and $a_l^*(x_0, x) = a_l(x)$ is this constant. If there exists i_0 such that $\partial a_l(x)/\partial x_{i_0}$ is not 0 then

$$a_l^*(x_0, x) = a_l(x) - x_0 \sum_{i=1}^m e_i \frac{\partial a_l(x)}{\partial x_i}$$

is invertible for all (x_0, x) with $x_0 \neq 0$. Hence the set U . ■

Remark. The hypothesis $|j| > |\alpha|$ is not an actual restriction.

THEOREM 4. Let $a_1, \dots, a_k, b_1, \dots, b_v$ be linear affine functions: $\mathbb{R}^m \rightarrow \mathbb{C}$. Let $a_{k+1}, \dots, a_{k+k'}, b_{v+1}, \dots, b_{v+v'}$ be linear affine functions, $\mathbb{R}^m \rightarrow \mathbb{R}$, each of the $a_1, \dots, a_{k+k'}$ non-identically zero. Let j, j', α, α' be multiindices such that $2|j| + |j'| > 2|\alpha| + |\alpha'|$ and $U = \{(x_0, x) \in \mathbb{R}^{m+1}; (|a_l(x_0, x)|^2)^* \text{ invertible for all } 1 \leq l \leq k \text{ and } a_l^*(x_0, x) \text{ invertible for all } k+1 \leq l' \leq k+k'\}$.

The Cauchy-Kovalewski extension of

$$\frac{|b_1(x)|^{2x_1} \dots |b_v(x)|^{2x_v} b_{v+1}^{x_{v+1}}(b)_{v+v'}^{x_{v+v'}}(x)}{|a_1(x)|^{2j_1} \dots |a_k(x)|^{2j_k} a_{k+1}^{j_1}(x) \dots a_{k+k'}^{j_{k'}}(x)}$$

is given for all $(x_0, x) \in U$ by

$$C_{2j+j', 2\alpha+\alpha'} D_{\zeta, 0}^{\alpha} \int_{A(x_0, x)} \frac{z^{j-1} \omega'(z)}{[\langle \zeta, b^*(x_0, x) \rangle + \langle z, a^*(x_0, x) \rangle]^{2|j|+|j'|-2|\alpha|-|\alpha'|}}$$

with

$$\begin{aligned} \langle \zeta, b^*(x_0, x) \rangle &= \sum_{l=1}^v \zeta_l b_l^*(x_0, x) + \sum_{l=1}^{v'} \zeta_{2v+l} b_{v+l}^*(x_0, x) \\ \langle z, a^*(x_0, x) \rangle &= \sum_{l=1}^k z_l a_l^*(x_0, x) + z_{k+1} \bar{a}_l^*(x_0, x) \\ &\quad + \sum_{l=1}^{k'} z_{2v+l} a_{k+l}^*(x_0, x). \end{aligned}$$

The harmonic extension of the previous rational function defined on \mathbb{R}^m is the real part of that Clifford-valued function.

Proof. By Proposition 6 we must show only $\{(x_0, x) \in \mathbb{R}^{m+1}; a_l^*(x_0, x) \text{ and } \bar{a}_l^*(x_0, x) \text{ invertible for all } 1 \leq l \leq k\} = \{(x_0, x) \in \mathbb{R}^{m+1}; (|a_l(x_0, x)|^2)^* \text{ invertible for all } 1 \leq l \leq k\}$.

Hence, we must show only

$$a_l^*(x_0, x) \text{ and } \bar{a}_l^*(x_0, x) \text{ invertible} \Leftrightarrow (|a_l(x_0, x)|^2)^* \text{ invertible.}$$

An element σ of a complexified Clifford algebra and its complex conjugate $\bar{\sigma}$ are such that σ and $\bar{\sigma}$ invertible $\Leftrightarrow \sigma \bar{\sigma}$ invertible; then

$$a_l^*(x_0, x) \text{ and } \bar{a}_l^*(x_0, x) \text{ invertible} \Leftrightarrow a_l^*(x_0, x) \bar{a}_l^*(x_0, x) \text{ invertible.}$$

But we must show that the extension of $|a_l(x)|^2$ is precisely that product:

$$\begin{aligned} (|a_l(x_0, x)|^2)^* &= a_l^*(x_0, x) \bar{a}_l^*(x_0, x) \\ a_l^*(x_0, x) \bar{a}_l^*(x_0, x) &= a_l \bar{a}_l - a_l x_0 \sum e_i \frac{\partial \bar{a}_l}{\partial x_i} - \bar{a}_l x_0 \sum e_i \frac{\partial a_l}{\partial x_i} \\ &\quad + x_0^2 \sum e_i e_i \frac{\partial a_l}{\partial x_i} \frac{\partial \bar{a}_l}{\partial x_i}. \end{aligned}$$

This is monogenic function on \mathbb{R}^{m+1} which is equal to $|a_l(x)|^2$ on \mathbb{R}^{m+1} . The Cauchy-Kovalewski extension is unique; then we get the inequality. ■

Remark. Intuitively the singularities do not see the non-commutativity structure.

3. A CONNECTION BETWEEN COMPLEX ANALYSIS AND CLIFFORD ANALYSIS

In [4], some connections between the two analyses were shown. Here we use the previous ideas to construct monogenic functions starting from holomorphic functions in a local manner.

Define the kernel

$$K(z, x_0, x) = (-1)^{k-1} (k-1)! \int_{A(z, x_0, x)} \frac{\omega'(\zeta)}{\langle \zeta, z - (x - ex_0) \rangle^m}$$

with

$$\langle \zeta, z - (x - ex_0) \rangle = \sum_{j=1}^m \zeta_j [z_j - (x_j - e_j x_0)].$$

$\Delta(z, x_0, x)$ is a simplex of the type defined in Theorem 1. This theorem shows that this kernel is defined for all z such that $z_j - (x_j - e_j x_0)$ is invertible; hence it is defined for all z such that $z_j \neq x_j \pm ix_0$.

Let $W = \{z \in \mathbb{C}^m; \operatorname{Im} z_1 = \pm \operatorname{Im} z_2 = \dots = \pm \operatorname{Im} z_m\}$. W is a 2^{m-1} covering of \mathbb{R}^{m+1} ,

$$(x_1 \pm ix_0, \dots, x_m \pm ix_0) \mapsto (x_0, x_1, \dots, x_m).$$

Then, we have the following.

PROPOSITION 7. *Let f be a holomorphic function defined in a neighborhood of W . Let*

$$P = \{z \in \mathbb{C}^m; |z - (x_j \pm ix_0)| < r_j \text{ for all } 1 \leq j \leq m\}$$

with r_j sufficiently small; then

$$F(x_0, x) = \int_{\partial P} f(z) K(z, x_0, x) d\sigma(z),$$

where ∂P is the distinguished boundary of P and $d\sigma$, the standard measure on it, is a monogenic function.

REFERENCES

1. F. BRACKX, R. DELANGHE, AND F. SOMMEN, "Clifford Analysis," Pitman Advanced Publishing, New York.
2. J. BURES AND V. SOUČEK, Generalised hypercomplex analysis and its integral formulas, *Complex Var.* **2** (1983), 139–150.
3. D. HESTENES AND G. SOBZYK, "Clifford Algebra to Geometric Calculus," Reidel Publishing Co., Dordrecht.
4. F. SOMMEN, Some connections between Clifford Analysis and complex analysis, *Complex Var.* **1** (1982), 97–118.